Vegetable garden:

The best producing cucumbers are burpless cucumbers when grown on a trellis.
 Cucumbers are heavy feeders. Keep them well fed with <u>Garden Trus</u>t. Be sure to grow burpless cumbers in a sunny location where the soil drains well.

Maple and Sycamore Trees:

- Maple and sycamore trees that are suffering from dying leaves likely have anthracnose. Treat by applying <u>Copper Fungicide</u> at bud break. Repeat at 7-10 day intervals during the spring.
- When you have a history of anthracnose in these trees, then make an application of *Copper Fungicide* at bud break and repeat at 7-10 day intervals twice to decrease the likelihood of future infections.
- Make sure these trees are fed with <u>Garden Trust</u>. When the weather gets warmer and dryer, these trees will have more energy to regrow new shoots and leaves.

Lawn:

Nutsedge is a pale green, fast growing, grass like plant that grows upright and faster
than bluegrass, tall fescue, or zoysia grass. Two days after mowing, nutsedge will be
clearly visible. It will be several inches taller than the rest of the lawn. Spray seedlings
of yellow nutsedge with Sedge hammer + as soon as you see nutsedge emerging in
your lawn. On southern lawns, they may already be visible. In the mid-Atlantic they will
be visible in early June. In Northern states they will be visible mid-summer.

Now is the time to apply your second application of <u>Turf Trust</u> to cool weather lawns. Zoysia lawns only need to be fed one time with Turf Trust per season. Feed your Zoysia lawn in late spring or early summer after it has turned fully green.

Zenith Zoysia in now available in seed and should be sown in the late spring or summer.

Shrubs:

- When planting azaleas or rhododendron, do not use bagged top soil, compost, or planting soils. Only use *Canadian Sphagnum Peat Moss* as a soil amendment.
- Never plant trees or shrubs deeper than the depth that they were in, in its previous container.
- English boxwoods turning rusty brown or orange are likely planted in wet, poorly draining soil. They are likely suffering from nematodes or phytophthora root rot. To treat the boxwood, start by testing the pH of the soil with a *Luster Leaf Ph tester*. The ideal pH is 6.7-7. When the pH is too low, use *Bonide Hydrate Lime*. If drainage/wet soil is a problem, then drill 18" deep holes with a 1.5" caliber earth auger. Make these holes 10" apart around the sick looking boxwood. Then fill these holes with Sharp Sand . Feed the boxwood with *Plant Trust Tree and Shrub Fertilizer*, but if *Bonide Hydrated Lime* was used, then wait to feed the boxwood for two weeks or for two heavy rains after the *Bonide Hydrated Lime* application. Continue to monitor the pH every six months. Avoid planing boxwoods in poorly draining soils. If you have wet soil, plant other plants that will better tolerate the wet soil.
- Hydrangeas will need extra attention this season, because of the late cold in the Northeast. Remove any dead cains and feed them with Garden Trust, as soon as

possible. Hydrangeas that where damaged by the late frost also will not bloom well this season.

Roses, Including Knockout roses, show holes in the leaves. Turn the leaves and you can see small 1/2 inch greenish leech like slugs underneath the rose leaves. These rose slugs cause the holes. Control rose slugs with Bonide eight spray and make sure that you feed the roses after treatment with Garden Trust 4 oz per plant.